

**Stony Brook University
The Graduate School**

Doctoral Defense Announcement

Abstract

Measurement of Direct Photons in Ultra-Relativistic Au+Au Collisions

By

Haijiang Gong

Direct photons provide a tool to study the different stages of a heavy ion collision, especially the formation of the quark-gluon-plasma (QGP), without being influenced by the strong reaction and hadronization processes. The yield of direct photons can be determined from the inclusive photon yield and the photon yield from hadronic decays. At low p_T , where a significant fraction of direct photon is expected to come from the thermalized medium of deconfined quarks, gluons and interacting hadrons, the measurement is very challenging. These so-called thermal photons carry information about the initial temperature of the medium

We present a new analysis technique that was developed to improve direct photon production measurement in the low and medium p_T range. The technique was applied to the PHENIX Run4 Au+Au $\sqrt{s_{NN}}=200\text{GeV}/c$ collisions. It uses strict particle identification (PID) in the Electromagnetic Calorimeter (EMCal) and a charged particle veto to extract a clean photon signal. These photons are then tagged with EMCal photon candidates with loose PID cuts, which can be reconstructed with high efficiency, to determine the fraction of photons originating from neutral pion decays. Most systematic uncertainties and detector effects cancel in this method. The results are compared with recent PHENIX direct photon measurement through external conversion method and theoretical calculation predicting thermal photon production.

Date: April 14th, 2014

Time: 2:30 AM

Place: Physics C-120

Program: Physics

Dissertation Advisor: Axel Drees